




APPENDIX A NUTRIENT MANAGEMENT PLAN TERMS (1 – 6) 1) LAND APPLICATION FIELD MAPS

UNITED FEEDERS LAND APPLICATION AREAS, WELLS WITHIN 0.5 MILES AND DISCHARGE PATH





Prepared by AgriTech Consulting
November 2012



Index to Soil Map Units

Map Abbreviation

Description

Soil Surface Texture (USDA)

NmA,B&C

Numa clay loam

Clay loam

RfA&B

Rocky Ford silty clay loam

Silty clay loam

SgC


Shingle loam


Clay loam

RgB

Rocky Ford silty clay loam, wet

Silty clay loam









APPENDIX B NUTRIENT MANAGEMENT PLAN TERMS 2) LAND APPLICATION INFORMATION





NMP TERMS - 2) LAND APPLICATION FIELDS

All land application fields are listed below.

Table B-1 – Land Application Fields

Field Identification	Latitude ¹	Longitude ²	Spreadable Acres ³
G-N	38.028622	-103.699865	61
G-S	38.699865	-103.699908	61
J-1	38.025242	-103.699007	30
J-2	38.021455	-103.69875	43
J-3	38.012354	-103.707461	37
-1	38.107973	-103.707161	46
-2	38.015437	-103.705702	28
-3	38.015708	-103.702312	35

¹Enter latitude in decimal degrees.

²Enter longitude in decimal degrees [number should be negative (eg. -104.3315)].

³Field acreages reduced by any setbacks, buffers, or otherwise unspreadable areas.





NMP TERMS - 2) LAND APPLICATION CROPS

All potential crops or other uses for each land application field are listed below.

Table -B-2 - Potential Land Application Field Crops

Field Identification	Crop	Realistic Yield Goal (See Appendix C)	Yield Unit (bushels, tons, etc.
G-N, G-S, U-1, U-2, U-3, L-1, L-2, L-3	Alfalfa	6.6	tons
G-N, G-S, U-1, U-2, U-3, L-1, L-2, L-3	Corn	220	Bushels
G-N, G-S, U-1, U-2, U-3, L-1, L-2, L-3	Grass	4.4	tons
G-N, G-S, U-1, U-2, U-3, L-1, L-2, L-3	Grain Sorghum	70.4	Bushels
G-N, G-S, U-1, U-2, U-3, L-1, L-2, L-3	Forage Sorghum	3.3	Tons
G-N, G-S, U-1, U-2, U-3, L-1, L-2, L-3	Sunflower	13.44	cwt
G-N, G-S, U-1, U-2, U-3, L-1, L-2, L-3	Winter Wheat	62.2	Bushels
G-N, G-S, U-1, U-2, U-3, L-1, L-2, L-3	Rye/Triticale	3.3	Tons
			,
			Section 1





NMP TERMS - 2) LAND APPLICATION CROP NUTRIENT NEEDS

Nutrient needs for each potential crop or other uses for each land application field are listed below.

Table B-3 - Crop Nutrient Needs

Crop	Yield Unit (bu, tons, etc.)	N Requirement (lbs/yield unit)	Information Source (see Appendix D)
Corn	Bu	*See attached page	CSUCE
Alfalfa	Tons	*See attached page	CSUCE
Grass	Tons	*See attached page	CSUCE
Winter Wheat	Bu	*See attached page	CSUCE
Grain Sorghum	Bu	*See attached page	NDSU
Forage Sorghum	Tons	*See attached page	NMSU
Sunflower	cwt	*See attached page	CSU, KSU, UN, WYU, USDA-ARS
Winter Wheat	Bu	*See attached page	CSUCE
Rye/Triticale	Tons	*See attached page	CSUCE
		1	
	1.		
		1	

NMP Terms- 2) Land Application Crop Nutrient Needs Table B- 3, N Requirements

**CSU recommendations are used, but are modified to account for the additional nitrogen that is mineralized form organic forms of nitrogen from previous applications. The percent of organic nitrogen that is mineralized from the second year it 10%, and the third year is 5%. These percentages are used for both liquid and solid nitrogen sources.

Alfalfa:

Nitrogen Rate calculated Using CSUCE publications:

"Update on Manure and Effluent Recommendations" no. 565
"Best Management Practices for Manure Utilization" Bulletin #XCM-174

- Gross N Need= Yield (tons) x 2000 lb x % Protein x 60% rate (N-fixing varieties) x 100 efficiency)

 Acre Ton 6.25
- N needed= Gross N Need -[Soil NO₃.N(lb/ac) in 0-24 in.] [%OM x 30(lb/ac)] [Irrigation Water NO₃-N (lb/ac)]
 Previous Crop or Manure Application N Credits (lbs/ac)]
- <u>N Needed</u> = Agronomic Rate (In/ac)
 Plant Available N

Corn:

Nitrogen Rate calculated Using CSUCE publications:

"Fertilizing Corn", no. 538

- N needed= 35 +[1.2 x EY(bu/ac)] -[Soil NO₃-N(lb/ac) in 0-24 in.] [%OM x 30(lb/ac)] [Irrigation Water NO₃-N (lb/ac)] [Previous Crop or Manure Application N Credits (lbs/ac)]
- <u>N Needed</u> = Agronomic Rate (In/ac)
 Plant Available N

Grasses:

Nitrogen Rate calculated Using CSUCE publications:

"Fertilizing Cool Season Grasses and Grass/legume Mixtures", no. 0.522

N needed= 45 lbs N x [EY (tons/acre) – 1.25 (tons/acre efficiency)] – [25 lbs/acre for every 6 ppm NO₃-N over 6 ppm.]- [Irrigation Water NO₃-N (lb/ac)]

Winter Wheat

Nitrogen Rate calculated Using CSUCE publications:

Gross N Need is determined using Tables 1 and 2 in CSU Fact Sheet no. 0.544

N is added according to CSU Publication if the wheat is grazed by cattle or managed to obtain protein levels greater than 12%.

- N needed= Gross N Need [Irrigation Water NO₃-N (lb/ac)] Previous Crop or Manure Application N Credits (lbs/ac)]
- <u>N Needed</u> = Agronomic Rate (In/ac)
 Plant Available N

Forage Sorghum

Nitrogen Rate calculated Using NMSU publications:

"Sorghum Forage Production in New Mexico", Guide A-332

- N needed= [8 x EY(bu/ac)] -[Soil NO₃.N(lb/ac) in 0-24 in.] [%OM x 30(lb/ac)] [Irrigation Water NO₃-N (lb/ac)]
 [Previous Crop or Manure Application N Credits (lbs/ac)]
- <u>N Needed</u> = Agronomic Rate (In/ac)
 Plant Available N

Rye/Triticale:

Nitrogen Rate calculated Using CSUCE publications:

"Best Management Practices for Manure Utilization" no. 568A

"Best Management Practices for Manure Utilization" Bulletin #XCM-174

- Gross N Need= <u>Yield (tons)</u> x <u>2000 lb</u> x 0.016 x <u>100</u> efficciency) Acre Ton 66
- N needed= Gross N Need -[Soil NO₃.N(lb/ac) in 0-24 in.] [%OM x 30(lb/ac)] [Irrigation Water NO₃-N (lb/ac)]
 Previous Crop or Manure Application N Credits (lbs/ac)]
- <u>N Needed</u> = Agronomic Rate (In/ac)
 Plant Available N

[&]quot;Fertilizing Winter Wheat", no. 0.544

[&]quot;Update on Manure and Effluent Recommendations" no. 565

[&]quot;Grain Protein Content and N Needs", no 0.555

Sunflower:

Nitrogen Rate calculated Using CSU, KSU, NU, WYU, USDA-ARS-Central Great Plains Research Station, Akron, Colorado, publication:

"High Plains Sunflower Production Handbook"

- N needed= [0.065 x EY(bu/ac) x STA] -[Soil NO₃₋N(lb/ac) in 0-24 in.] [%OM x 30(lb/ac)] [Irrigation Water NO₃-N (lb/ac)] [Previous Crop or Manure Application N Credits (lbs/ac)]
- STA= Soil Texture Adjustment (1.1 for sandy soils less than 1.0 percent OM, 1.0 for other soils)
- N Needed = Agronomic Rate (In/ac)
 Plant Available N

Grain Sorghum (Milo):

Nitrogen Rate calculated Using NDSU publications:

"Grain Sorghum (Milo) Production Guidelines"

- N needed= [1.1 x EY(bu/ac)] -[Soil NO₃-N(lb/ac) in 0-24 in.] [%OM x 30(lb/ac)] [Irrigation Water NO₃-N (lb/ac)] [Previous Crop or Manure Application N Credits (lbs/ac)]
- <u>N Needed</u> = Agronomic Rate (In/ac)
 Plant Available N





APPENDIX C

NUTRIENT MANAGEMENT PLAN TERMS

3) EXPECTED CROP YIELD INFORMATION





Historical crop yield information source: http://www.nass.usda.gov/Data_and_Statistics/Quick_Stats_1.0/index.asp

Facility Name: United		United	Feeders
Field Iden	tification:	G-N, G-S, U-1,	L-1
Crop:		Corn	
Column A	Column B		
Year	Yield	Units (bu/ac, tons, etc.)	Notes: (i.e. drought, flood)
2008	210	Bu/acre	
2009	210	Bu/acre	
2010	200	Bu/acre	drought
2011	200	Bu/acre	drought
2012	200	Bu/acre	drought
TOTAL:	Total Bushels (Sum of Column B)	/ 5 # of Years (from Column A)	= <u>204</u> +10% <u>220.4</u> Average Realistic Yield Goal

REALISTIC YIELD GOAL WORKSHEET

Historical crop yield information source: http://www.nass.usda.gov/Data and Statistics/Quick Stats 1.0/index.asp

Facility N	ame:	United Feeders	
Field Iden	tification:	<u>L2</u>	
Crop:		Grass Hay	and the second s
Column A	Column B		
Year	Yield	Units (bu/ac, tons, etc.)	Notes: (i.e. drought, flood)
2010	4	Tons/acre	Drought
2011	4	Tons/acre	Drought
TOTAL:	The second of the	8 /	2 =4 +10% _4.4
	Total Bushels (Sum of Column B)	# of Years (from Column A)	Average Realistic Yield Goal





Historical crop yield information source: http://www.nass.usda.gov/Data_and_Statistics/Quick_Stats_1.0/index.asp

Facility N	ame:	United	Feeders
Field Iden	tification:	U-3	
Crop:		Alfalfa	
Column A	Column B		
Year	Yield	Units (bu/ac, tons, etc.)	Notes: (i.e. drought, flood)
2008	6	Tons/acre	Drought
2009	6	Tons/acre	Drought
2010	6	Tons/acre	Drought
TOTAL:	18	_ /	<u>3</u> = <u>6</u> +10% <u>66</u>
	Total Bushels (Sum of Column B)	# of Years (from Column A)	Average Realistic Yield Goal

REALISTIC YIELD GOAL WORKSHEET

Historical crop yield information source: http://www.nass.usda.gov/Data_and_Statistics/Quick_Stats_1.0/index.asp

Facility N	ame:	United Fe	eders
Field Iden	tification:	All	
Crop:		Grain Sorghum	
Column A	Column B		
Year	Yield	Units (bu/ac, tons, etc.)	Notes: (i.e. drought, flood)
2010	64	Bu/acre	Colorado Ag. Statistics, Drought
TOTAL:	Total Bushels (Sum of Column	/ 1 # of Years	Average Realistic Yield Goal





Historical crop yield information source: http://www.nass.usda.gov/Data_and_Statistics/Quick_Stats_1.0/index.asp

Facility N	ame:	United	Feeders
Field Iden	tification:	All	
Crop:		Forage Sorghum	1
Column A	Column B		
Year	Yield	Units (bu/ac, tons, etc.)	Notes: (i.e. drought, flood)
2006-11	3	Ton/acre	USDA-FSA County Avg., Drought
TOTAL:	15	_ / 5	= 3 +10% 3.3
	Total Bushels (Sum of Column B)	# of Years (from Column A)	Average Realistic Yield Goal

REALISTIC YIELD GOAL WORKSHEET

Historical crop yield information source: http://www.nass.usda.gov/Data_and_Statistics/Quick_Stats_1.0/index.asp

Facility Name:		United Feed	lers
Field Iden	tification:	All	
Crop:		Sunflower	
Column A	Column B		
Year	Yield	Units (bu/ac, tons, etc.)	Notes: (i.e. drought, flood)
2006-11	12.22	Cwt/acre	USDA-FSA County Avg., Drought
TOTAL:	Total Bushels (Sum of Column I	/ <u>5</u> # of Years B) (from Colu	Average Realistic Yield Goal





Historical crop yield information source: http://www.nass.usda.gov/Data and Statistics/Quick Stats 1.0/index.asp

Facility Name:		United	Feeders
Field Iden	tification:	All	
Crop:		Winter Wheat	
Column A	Column B		
Year	Yield	Units (bu/ac, tons, etc.)	Notes: (i.e. drought, flood)
2010	56.5	Bu/acre	Colorado Ag Statistics, Drought
TOTAL:	56.5	/ 1	= 56.5 +10% 62.2
	Total Bushels (Sum of Column B)	# of Years (from Column A)	Average Realistic Yield Goal

REALISTIC YIELD GOAL WORKSHEET

Historical crop yield information source: http://www.nass.usda.gov/Data_and_Statistics/Quick_Stats_1.0/index.asp

Facility Name: Field Identification:		United Feeder	S
Crop:	tification:	All Rye/Triticale	
Column A	Column B		
Year	Yield	Units (bu/ac, tons, etc.)	Notes: (i.e. drought, flood)
2006-11	3	Tons/acre	USDA-FSA County Avg., Drought
TOTAL:	15	/ 5	= <u>3</u> +10% <u>3.3</u>
	Total Bushels (Sum of Column B)	# of years (From Col	





APPENDIX D NUTRIENT MANAGEMENT PLAN TERMS 4) NUTRIENT BUDGET INFORMATION





4) NUTRIENT BUDGET INFORMATION

Nutrient Budge					
Crop:	Manure and Process Wastewater Application Rate Calculated Using:	Description of Method to be Used (calculation, look-up table, etc.):			
Alfalfa	☐ CSUCE Published Fertilizer Suggestions ☐ Adjacent State CE-Published Fertilizer Suggestions ☐ CNMP Method that meets USDA-NRCS standards ☐ CO NRCS NMP guidelines ☐ Ag Program-approved Method	"Update on Manure and Effluent Reccomendations", CSUCE no. 565; "Best Management Practices for Manure Utilization", CSU Bulletin #XCM-174			
<u>Corn</u>	X CSUCE Published Fertilizer Suggestions Adjacent State CE-Published Fertilizer Suggestions CNMP Method that meets USDA-NRCS standards CO NRCS NMP guidelines Ag Program-approved Method	"Fertilizing Corn", CSUCE Fact Sheet no. 0.538, (10/09)			
Grain Sorghum (Milo)	☐ CSUCE Published Fertilizer Suggestions ☐ Adjacent State CE-Published Fertilizer Suggestions ☐ CNMP Method that meets USDA-NRCS standards ☐ CO NRCS NMP guidelines ☐ Ag Program-approved Method	"Grain Sorghum (Milo) Production Guidelines", NDSU			
Forage Sorghum	☐ CSUCE Published Fertilizer Suggestions ☐ Adjacent State CE-Published Fertilizer Suggestions CMMP Method that meets USDA-NRCS standards CO NRCS NMP guidelines ☐ Ag Program-approved Method	"Sorghum Forage Production in New Mexico", NMSU Guide A-332, (8/11)			
Rye <u>/Tritic</u> ale	 ∑ CSUCE Published Fertilizer Suggestions ☐ Adjacent State CE-Published Fertilizer Suggestions ☐ CNMP Method that meets USDA-NRCS standards ☐ CO NRCS NMP guidelines ☐ Ag Program-approved Method 	"Best Management Practices for Manure Utilization", CSUCE no. 568A; "Best Management Practices for Manure Utilization", CSU Bulletin #XCM-174			
Winter Wheat	 \(\times \) CSUCE Published Fertilizer Suggestions \(\times \) Adjacent State CE-Published Fertilizer Suggestions \(\times \) CNMP Method that meets USDA-NRCS standards \(\times \) CO NRCS NMP guidelines \(\times \) Ag Program-approved Method 	"Fertilizing Winter Wheat", CSUCE Fact Sheet no. 0.544 (5/09); "Grain Protein Content and N needs", CSUCE Fact Sheet no. 0.555, (1/08)			
Su <u>nflow</u> er	☐ CSUCE Published Fertilizer Suggestions ☐ Adjacent State CE-Published Fertilizer Suggestions ☐ CNMP Method that meets USDA-NRCS standards ☐ CO NRCS NMP guidelines ☐ Ag Program-approved Method	"High Plains Sunflower Production Handbook", CSU, KSU, UN, WYU, USDA- ARS-Central Great Plains Research Station, Akron, CO.			
Wh <u>eatgra</u> ss, Bromegrass, Fescue	 \(\times \) CSUCE Published Fertilizer Suggestions \(\times \) Adjacent State CE-Published Fertilizer Suggestions \(\times \) CNMP Method that meets USDA-NRCS standards \(\times \) CO NRCS NMP guidelines \(\times \) Ag Program-approved Method 	"Fertilizing Cool Season Grasses and Grass/Legume Mixtures", CSUCE Fact Sheet no. 0.522, (1/11)			
Realistic yield go	oals determined using worksheet(s) is Appendix C? [X	☐ Yes ☐ No			
Realistic yield goals determined using methods other than worksheet(s) is Appendix C? Yes X No If yes, describe how realistic yield goals will be determined (crop insurance factors should not be added to yield goals): Colorado Ag. Statistics figures and County averages if no history is available					





APPENDIX E

NUTRIENT MANAGEMENT PLAN TERMS

5) COLORADO PHOSPHORUS INDEX RISK ASSESSMENT






5) PHOSPHORUS AND NITROGEN TRANSPORT

Application rates for manure and process wastewater applied to land application sites minimize phosphorus and nitrogen transport from the application sites to surface waters. An initial assessment of the potential for phosphorus and nitrogen transport risk to surface water will be made prior to manure or process wastewater being applied to an application site. [Regulation No. 61.17(8)(b)(xii)(B)]

There is currently no published tool suitable for assessing nitrogen transport risk. Phosphorus and nitrogen transport risk will be assessed using the Colorado Phosphorus Index Risk Assessment.

The following flow chart will be used to determine if a phosphorus risk assessment must be completed for a land application site:







5) PHOSPHORUS AND NITROGEN TRANSPORT (continued)

For land application fields that require a Colorado Phosphorus Index Risk Assessment to be completed, the following applicable best management practices will be incorporated:

- (A) Phosphorus-based manure and process wastewater application rates may be made to application sites where the risk of off-site phosphorus transport is scored as high.
- (B) No application of manure or process wastewater will be made to land application sites where the risk of off-site phosphorus transport is rated as very high¹.
- (C) No application of manure or process wastewater will be made to a land application site where the risk of off-site nitrogen transport to surface water is not minimized.
- (D) Where a multi-year phosphorus application was made to a land application site, no additional manure or process wastewater will be applied to the same site in subsequent years until the applied phosphorus has been removed from the site via harvest and crop removal.

After completing an initial assessment of the potential for phosphorus and/or nitrogen transport from a land application site to surface water, additional assessments will be made every five years or at the frequency described below, whichever is sooner:

Cause for Re-Assessment	Frequency
Where a crop management change has occurred	For phosphorus - Assess within one year after such a change would reasonably result in an increase in the transport risk assessment score. For nitrogen – Assess within one year after such a change would reasonably result in the nitrogen transport to surface water not being minimized.
Where a phosphorus transport risk assessment score was very high	Assess phosphorus transport risk within six months of intending to apply manure or process wastewater, except where the initial assessment is scored as very high, then there shall be a three-year period within which to manage the site for the purpose of lowering the phosphorus transport risk assessment rating to high or less. During this period, manure or process wastewater may be applied to the site at either nitrogen- or phosphorus-based rates.
Where a nitrogen transport risk assessment reveals that nitrogen transport to surface water is not minimized	Assess nitrogen transport risk within six months of intending to apply manure or process wastewater.

ASSOCIATED RECORDS:

1) Copies of phosphorus/nitrogen transport risk assessments are maintained on-site.

Where the initial assessment of a land application site scores very high, the facility has a three-year period within which to manage the site for the purpose of lowering the phosphorus transport risk assessment rating to high or lower. During this period, manure or process wastewater may be applied to the site at either nitrogen- or phosphorus-based rates.

Cooperator	United Feeders	Crop Year	2012	Date	7-29-12
Tract//Field No	G-N & S	Acres	122	Planner	M. Nelson
Previous Crop	Corn	Planned Crop	Corn	Yield Goal	200 Bu/Ac.

Factor\Ris	k	Low (1)	Medium (2)	High (3)	Very High (4)	Score
1a. Runoff Clas (Irrigated Sites Onl Table 1a) OR		Negligible, Very Low or Low	Medium	High	Very High	1
1b. Rill and Inte Erosion (Non-Imig Sites Only - See Ta	gated	<3	3-5	6-10	> 10	
2. Soil Test P (S Table 2)	See	Low	Medium	High	Very High	4
3. P ₂ O ₅ Applica Rate (See Table 3a. An applied or rotations	nually	< 30	30-90	91-150	> 150	1
4. P Application Method (See Table 4. Use highest applicable risk category for multiple P applications)		P is not applied, or P is injected or subsurface applied	Spring applied and incorporated within 2 weeks, or sprinkler applied in-season	Fall/winter applied and incorporated within 2 weeks, and fall topdress for surface or flood irrigated pasture or hayland	Surface applied w/ no incorporation, and spring topdress for surface or flood irrigated pasture or hayland	2
Gross Score (S	um of l	Factors 1 throug	h 4)			. 8
5. BMP, Subtract one point for each of the following BMPs or Mitigation Factors applicable to this site. Contour Buffer Strips, Cover Crops, Filter Strips, Furrow Diking, Grassed Waterways, Linear Polyacrylamide, Terraces, Soil Test Free Lime > 4%, or Residue and Tillage Management						-1
Not Score (Sun	n of Fac	tors 1 through 4	Liess Factor 5 RMP in	nplementation Credits)		7

Net Score	Risk Interpretations
< 8	This field has a LOW potential for off-site P movement if managed at the current level. Calculate organic nutrient application rates according to crop nitrogen requirements.
8 to 11	This field has a MEDIUM potential for off-site P movement. Consider management changes to decrease risk and support continued long-term organic nutrient applications. Calculate organic nutrient application rates according to crop nitrogen requirements.
12 to 15	This field has a HIGH potential for off-site P movement. Implement management changes to decrease risk. Calculate organic nutrient application rates according to crop phosphorus requirements.
16	This field has a VERY HIGH potential for off-site P movement. Implement management changes to decrease risk. Do not apply organic nutrients to this field without decreasing the risk for off-site transport

Cooperator	United Feeders	Crop Year	2012	Date	7-29-12
Tract//Field No	U-1	Acres	30	Planner	M. Nelson
Previous Crop	Corn	Planned Crop	Corn	Yield Goal	200 Bu/Ac.

Factor\Ris	k	Low (1)	Medium (2)	High (3)	Very High (4)	Score
1a. Runoff Clas (Irrigated Sites Onl Table 1a) OR		Negligible, Very Low or Low	Medium	High	Very High	1
1b. Rill and Inte Erosion (Non-Irri Sites Only - See T	gated	<3	3-5	6-10	> 10	
2. Soil Test P (S Table 2)	See	Low	Medium	High	Very High	2
3. P ₂ O ₅ Applica Rate (See Table 3a. An applied or rotations	nually	< 30	30-90	91-150	> 150	1
4. P Application Method (See Table 4. Use highest applicable risk category for multiple P applications)		P is not applied, or P is injected or subsurface applied	Spring applied and incorporated within 2 weeks, or sprinkler applied in-season	Fall/winter applied and incorporated within 2 weeks, and fall topdress for surface or flood irrigated pasture or hayland	Surface applied w/ no incorporation, and spring topdress for surface or flood irrigated pasture or hayland	4
Gross Score (S	Sum of I	Factors 1 throug	h 4)	I		. 8
5. BMP, Mitigation Factor Credits	Contou	r Buffer Strips, C	over Crops, Filter Strips	s or Mitigation Factors ap s, Furrow Diking, Grasse > 4%, or Residue and Till	d Waterways, Linear	-1
Net Score (Sur	n of Fac	etors 1 through 4	l less Factor 5, BMP in	polementation Credits)		7

Net Score	Risk Interpretations
< 8	This field has a LOW potential for off-site P movement if managed at the current level. Calculate organic nutrient application rates according to crop nitrogen requirements.
8 to 11	This field has a MEDIUM potential for off-site P movement. Consider management changes to decrease risk and support continued long-term organic nutrient applications. Calculate organic nutrient application rates according to crop nitrogen requirements.
12 to 15	This field has a HIGH potential for off-site P movement. Implement management changes to decrease risk. Calculate organic nutrient application rates according to crop phosphorus requirements.
16	This field has a VERY HIGH potential for off-site P movement. Implement management changes to decrease risk. Do not apply organic nutrients to this field without decreasing the risk for off-site transport

Cooperator	United Feeders	Crop Year	2012	Date	7-29-12
Tract//Field No	U-3	Acres	37	Planner	M. Nelson
Previous Crop	Corn	Planned Crop	Corn	Yield Goal	200 Bu/Ac.

Factor\Risk	(Low (1)	Medium (2)	High (3)	Very High (4)	Score
1a. Runoff Class (Irrigated Sites Only - See Table 1a) OR		Negligible, Very Low or Low	Medium	. High	Very High	1
1b. Rill and Inter Erosion (Non-Irrig Sites Only - See Ta	ated	<3	3-5	6-10	> 10	
2. Soil Test P (Se Table 2)	ee	Low	Medium	High	Very High	3
 P₂O₅ Applicat Rate (See Table 3a. Annapplied or rotational 	nually	< 30	30-90	91-150	> 150	1
4. P Application Method (See Table 4. Use highest applicable risk category for multiple P applications)		P is not applied, or P is injected or subsurface applied	Spring applied and incorporated within 2 weeks, or sprinkler applied in-season	Fall/winter applied and incorporated within 2 weeks, and fall topdress for surface or flood irrigated pasture or hayland	Surface applied w/ no incorporation, and spring topdress for surface or flood irrigated pasture or hayland	4
Gross Score (Si	um of F	Factors 1 throug	h 4)			. 9
5. BMP, Mitigation Factor Credits Subtract one point for each of the following BMPs or Mitigation Factors applicable to this site. Contour Buffer Strips, Cover Crops, Filter Strips, Furrow Diking, Grassed Waterways, Linear Polyacrylamide, Terraces, Soil Test Free Lime > 4%, or Residue and Tillage Management					-1	
	122.000			nplementation Credits)		8

Net Score	Risk Interpretations
< 8	This field has a LOW potential for off-site P movement if managed at the current level. Calculate organic nutrient application rates according to crop nitrogen requirements.
8 to 11	This field has a MEDIUM potential for off-site P movement. Consider management changes to decrease risk and support continued long-term organic nutrient applications. Calculate organic nutrient application rates according to crop nitrogen requirements.
12 to 15	This field has a HIGH potential for off-site P movement. Implement management changes to decrease risk. Calculate organic nutrient application rates according to crop phosphorus requirements.
16	This field has a VERY HIGH potential for off-site P movement. Implement management changes to decrease risk. Do not apply organic nutrients to this field without decreasing the risk for off-site transpor

Cooperator	United Feeders	Crop Year	2012	Date	7-29-12
Tract//Field No	L-1	Acres	46	Planner	M. Nelson
Previous Crop	Corn	Planned Crop	Corn	Yield Goal	200 Bu/Ac.

Factor\Ris	k	Low (1)	Medium (2)	High (3)	Very High (4)	Score
1a. Runoff Class (Irrigated Sites Only - See Table 1a) OR		Negligible, Very Low or Low	Medium	High	Very High	1
1b. Rill and Inte Erosion (Non-Irrig Sites Only - See To	gated	<3	3-5	6-10	> 10	
2. Soil Test P (S Table 2)	See	Low	Medium	High	Very High	3
3. P ₂ O ₅ Applica Rate (See Table 3a. An applied or rotations	nually	< 30	30-90	91-150	> 150	1
4. P Application Method (See Table 4. Use highest applicable risk category for multiple P applications)		P is not applied, or P is injected or subsurface applied	Spring applied and incorporated within 2 weeks, or sprinkler applied in-season	Fall/winter applied and incorporated within 2 weeks, and fall topdress for surface or flood irrigated pasture or hayland	Surface applied w/ no incorporation, and spring topdress for surface or flood irrigated pasture or hayland	4
Gross Score (S	um of I	actors 1 throug	h 4)			9
5. BMP, Mitigation Factor Credits Subtract one point for each of the following BMPs or Mitigation Factors applicable to this site. Contour Buffer Strips, Cover Crops, Filter Strips, Furrow Diking, Grassed Waterways, Linear Polyacrylamide, Terraces, Soil Test Free Lime > 4%, or Residue and Tillage Management					-1	
Net Score (Sun	n of Fac	tors 1 through 4	l less Factor 5, BMP In	nplementation Credits)		8

Net Score	Risk Interpretations
< 8	This field has a LOW potential for off-site P movement if managed at the current level. Calculate organic nutrient application rates according to crop nitrogen requirements.
8 to 11	This field has a MEDIUM potential for off-site P movement. Consider management changes to decrease risk and support continued long-term organic nutrient applications. Calculate organic nutrient application rates according to crop nitrogen requirements.
12 to 15	This field has a HIGH potential for off-site P movement. Implement management changes to decrease risk. Calculate organic nutrient application rates according to crop phosphorus requirements.
16	This field has a VERY HIGH potential for off-site P movement. Implement management changes to decrease risk. Do not apply organic nutrients to this field without decreasing the risk for off-site transport

Cooperator	United Feeders	Crop Year	2012	Date	7-29-12	
Tract//Field No	L-3	Acres	35	Planner	M. Nelson	
Previous Crop	Corn	Planned Crop	Fallow	Yield Goal	0	

Factor\Ris	sk	Low (1)	Medium (2)	High (3)	Very High (4)	Score
1a. Runoff Clas (Irrigated Sites On Table 1a) OR		Negligible, Very Low or Low	Medium	High	Very High	1
1b. Rill and Inte Erosion (Non-Im Sites Only - See T	igated	<3	3-5	6-10	> 10	
2. Soil Test P (Table 2)	See	Low	Medium	High	Very High	3
3. P ₂ O ₅ Applica Rate (See Table 3a. An applied or rotation	nnually	< 30	30-90	91-150	> 150	1
4. P Application Method (See Table 4. Uhighest applicate category for mulapplications)	lse ole risk	P is not applied, or P is injected or subsurface applied	Spring applied and incorporated within 2 weeks, or sprinkler applied in-season	Fall/winter applied and incorporated within 2 weeks, and fall topdress for surface or flood irrigated pasture or hayland	Surface applied w/ no incorporation, and spring topdress for surface or flood irrigated pasture or hayland	4
Gross Score (Sum of I	Factors 1 throug	h 4)		L .	. 9
5. BMP, Mitigation Factor Credits	Contou	Buffer Strips, C	over Crops, Filter Strips	s or Mitigation Factors app s, Furrow Diking, Grasse > 4%, or Residue and Till	d Waterways, Linear	-1
Not Score (Sur	n of Ear	store 4 through	l less Factor 5, BMP In			8

Net Score	Risk Interpretations
<8	This field has a LOW potential for off-site P movement if managed at the current level. Calculate organic nutrient application rates according to crop nitrogen requirements.
8 to 11	This field has a MEDIUM potential for off-site P movement. Consider management changes to decrease risk and support continued long-term organic nutrient applications. Calculate organic nutrient application rates according to crop nitrogen requirements.
12 to 15	This field has a HIGH potential for off-site P movement. Implement management changes to decrease risk. Calculate organic nutrient application rates according to crop phosphorus requirements.
16	This field has a VERY HIGH potential for off-site P movement. Implement management changes to decrease risk. Do not apply organic nutrients to this field without decreasing the risk for off-site transport





APPENDIX F NUTRIENT MANAGEMENT PLAN TERMS 6) FIELD NUTRIENT BALANCE CALCULATIONS





(Conduct calculations for each crop, for each field)

Facility Name: United Feeders, Inc.

PermitNumber: 931038

Land Application Site Name: G-N & G-S (also known as G1

Date: 7-29-12

Table F-1

	(crop sequence/ro	tation and year (circle current cr	op)	Realistic Yield Goal
Year	2012	2013	2014	2015	2016	(of current year)
Crop	corn	corn	corn	corn	corn	220 bu/ac

Table F-2

Current soil	test levels	(ppm or lb/	ac)				
Soil Test Date	N* (as NO ₃ -N)	P*	Phosphorus Test Extraction Used (AB-DTPA, Bray, Mehlich, NaHCO ₃)	K	pН	CEC	O.M.%
2/29/12	97 lbs	100 ppm	NaHCO3	459	8.0	.84	2.1

^{*}Must be tested

Table F-3

Recommend	ed nutrients/amendme	ents to meet realistic yi	ield goal (see Appendi	x B, Table B-3)
N^1	P ₂ O ₅	K ₂ O	Lime	Other:
299 lbs	0	0		

N number is based on removal, rather than on soil test NO₃-N carryover from the previous crop. Use this value to complete line 10 on Table F-4 below.

Table F-4

	Nutrient Sources	N
		²lbs/ac
1. Nitrogen cre	dits from previous legume crop	0
2. Nitrogen cre	dit from irrigation water	Na
3. Other (e.g., s	oil organic matter mineralization, atmospheric deposition/evaporation)	63
4. Soil nitroger	n credit	97
5.	Total credits	160
		N
6. Credits (from	n row 5 above)	160
7. Plant availal	ole nitrogen (PAN) content of manure, litter, and process wastewater	0 - no app in 2012
8. Fertilizer	Starter	0
	Other	139
9.	Subtotal (sum of line 6, 7, and 8)	299
10.	Nitrogen recommended (from Table F-3)	299
11.	Nitrogen Status (subtract line 10 from line 9)	0

²Use the same units for each line in Table F-4. Include documentation of unit conversion factors used, if any.

	Nutrient Management Specifications				
Amount to be applied (lb/ac)	N:139	P ₂ O ₅ :0	K ₂ O:0	Other:0	

Predicted method, form, and timing of application: Split fertigation applications and/or sidedress in June





(Conduct calculations for each crop, for each field)

Facility Name: United Feeders, Inc.

PermitNumber: 931038

Land Application Site Name: <u>U1</u>

Date: 7-29-12

Table F-1

	(crop sequence/ro	tation and year (circle current cr	op)	Realistic Yield Goal
Year	2012	2013	2014	2015	2016	(of current year)
Crop	corn	corn	corn	corn	corn	220 Bu

Table F-2

Current soil	test levels	(ppm or lb.	/ac)				
Soil Test Date	N* (as NO ₃ -N)	P*	Phosphorus Test Extraction Used (AB-DTPA, Bray, Mehlich, NaHCO ₃)	К	pН	CEC	O.M.%
2-27-12	111 lbs	35 ppm	NaHCO3	236 ppm	8.0	1.94	1.7

^{*}Must be tested

Table F-3

Recommend	ed nutrients/amendme	nts to meet realistic yi	ield goal (see Appendi	x B, Table B-3)
N^1	P_2O_5	K ₂ O	Lime	Other:
299 lbs	20 lbs	0	1 <u>-1-1</u>	

N number is based on removal, rather than on soil test NO₃-N carryover from the previous crop. Use this value to complete line 10 on Table F-4 below.

Table F-4

	N	
		²lbs/ac
1. Nitrogen cre	dits from previous legume crop	0
2. Nitrogen cre	dit from irrigation water	Na
3. Other (e.g., s	oil organic matter mineralization, atmospheric deposition/evaporation)	51
4. Soil nitroger	credit	111
5.	Total credits	162
		N
6. Credits (from	n row 5 above)	162
7. Plant availal	ole nitrogen (PAN) content of manure, litter, and process wastewater	0- no app in 2012
8. Fertilizer	Starter	0
o. Termizer	Other	137
9.	Subtotal (sum of line 6, 7, and 8)	299
	Nitrogen recommended (from Table F-3)	299
10.		0

²Use the same units for each line in Table F-4. Include documentation of unit conversion factors used, if any.

Nutrient Management Specifications						
Amount to be applied (lb/ac)	N:137	P ₂ O ₅ :20	K2O:0	Other:0		

Predicted method, form, and timing of application: Sidedress crop in early June





(Conduct calculations for each crop, for each field)

Facility Name: <u>United Feeders. Inc.</u> PermitNumber: <u>931038</u>

Land Application Site Name: <u>U-3</u> Date: <u>7-29-12</u>

Table F-1

	Cro	Realistic Yield Goal				
Year	2012	2013	2014	2015	2016	(of current year)
Crop	New alfalfa	alfalfa	alfalfa	alfalfa	alfalfa	4 tons/acre

Table F-2

Current soil test levels (ppm or lb/ac)							
Soil Test Date	N* (as NO ₃ -N)	P*	Phosphorus Test Extraction Used (AB-DTPA, Bray, Mehlich, NaHCO ₃)	K	pН	CEC	O.M.%
2/27/12	149 lbs.	59 ppm	NaHCO3	301 ppm	7.9	2.04	2.0

^{*}Must be tested

Table F-3

Recommende	d nutrients/amendme	ents to meet realistic yi	ield goal (see Appendi	x B, Table B-3)
N^1	P_2O_5	K ₂ O	Lime	Other:
0 lbs/acre	0	0		

N number is based on removal, rather than on soil test NO₃-N carryover from the previous crop. Use this value to complete line 10 on Table F-4 below.

Table F-4

	Nutrient Sources	N
		²lbs/ac
1. Nitrogen cre	dits from previous legume crop	90
2. Nitrogen cre	dit from irrigation water	na
3. Other (e.g., s	oil organic matter mineralization, atmospheric deposition/evaporation)	60
4. Soil nitroger	n credit	149
5.	Total credits	299
		N
6. Credits (from	n row 5 above)	299
7. Plant availal	ble nitrogen (PAN) content of manure, litter, and process wastewater	0- no app in 2012
8. Fertilizer	Starter	0
0. 10.000	Other	0
9.	Subtotal (sum of line 6, 7, and 8)	299
10.	Nitrogen recommended (from Table F-3)	210
11.	Nitrogen Status (subtract line 10 from line 9)	89

²Use the same units for each line in Table F-4. Include documentation of unit conversion factors used, if any.

Nutrient Management Specifications					
Amount to be applied (lb/ac)	N:0	P ₂ O ₅ :0	K ₂ O:0	Other:0	

Predicted method, form, and timing of application: None





(Conduct calculations for each crop, for each field)

Facility Name: United Feeders. Inc.

PermitNumber: 931038

Land Application Site Name: L-1

Date: 7-29-12

Table F-1

	(Realistic Yield Goal				
Year	2012	2013	2014	2015	2016	(of current year)
Crop	corn	corn	corn	corn	corn	220 bu

Table F-2

Current soil test levels (ppm or lb/ac)							
Soil Test Date	N* (as NO ₃ -N)	P*	Phosphorus Test Extraction Used (AB-DTPA, Bray, Mehlich, NaHCO ₃)	K	pН	CEC	O.M.%
2-29-12	69 lbs	44 ppm	NaHCO3	384 ppm	7.8	3.84	1.9

^{*}Must be tested

Table F-3

Recommend	ded nutrients/amendme	ents to meet realistic yi	ield goal (see Appendi	x B, Table B-3)
N^1	P ₂ O ₅	K ₂ O	Lime	Other:
299	0	0	0	0

N number is based on removal, rather than on soil test NO₃-N carryover from the previous crop. Use this value to complete line 10 on Table F-4 below.

Table F-4

	Nutrient Sources					
		²lbs/ac				
1. Nitrogen cre	dits from previous legume crop	0				
2. Nitrogen cre	dit from irrigation water	na				
3. Other (e.g., s	oil organic matter mineralization, atmospheric deposition/evaporation)	57				
4. Soil nitroger	credit	69				
5.	Total credits	126				
		N				
6. Credits (from	n row 5 above)	126				
7. Plant availab	ole nitrogen (PAN) content of manure, litter, and process wastewater	0- no app in 2012				
8. Fertilizer	Starter	0				
	Other	173				
9.	Subtotal (sum of line 6, 7, and 8)	299				
10.	Nitrogen recommended (from Table F-3)	299				
11.	Nitrogen Status (subtract line 10 from line 9)	0				

²Use the same units for each line in Table F-4. Include documentation of unit conversion factors used, if any.

Nutrient Management Specifications						
Amount to be applied (lb/ac)	N:173	P ₂ O ₅ :0	K ₂ O:0	Other:0		

Predicted method, form, and timing of application: Sidedress crop in June





(Conduct calculations for each crop, for each field)

Facility Name: United Feeders, Inc.

PermitNumber: 931038

Land Application Site Name: L-3

Date: 7-29-12

Table F-1

Crop sequence/rotation and year (circle current crop)						Realistic Yield Goal	
Year	2012	2013	2014	2015	2016	(of current year)	
Crop	fallow	corn	corn	fallow	corn	220 Bu/ac	

Table F-2

Current soil	test levels	(ppm or lb.	/ac)				
Soil Test Date	N* (as NO ₃ -N)	P*	Phosphorus Test Extraction Used (AB-DTPA, Bray, Mehlich, NaHCO ₃)	K	pН	CEC	O.M.%
2-22-12	135 lbs	49 ppm	NaHCO3	373 ppm	7.9	2.68	2.4

^{*}Must be tested

Table F-3

Recommend	ded nutrients/amendme	ents to meet realistic yi	ield goal (see Appendi	x B, Table B-3)
N^1	P ₂ O ₅	K ₂ O	Lime	Other:
0	0	0	0	0

N number is based on removal, rather than on soil test NO₃-N carryover from the previous crop. Use this value to complete line 10 on Table F-4 below.

Table F-4

1 Nitrogon aradi		²lbs/ac	
1 Nitrogen andi		103/40	
1. Millogen credi	its from previous legume crop	0	
2. Nitrogen credi	Na		
3. Other (e.g., soi	72		
4. Soil nitrogen of	credit	135	
5.			
		N	
6. Credits (from	row 5 above)	207	
7. Plant available	e nitrogen (PAN) content of manure, litter, and process wastewater	0- no app in 2012	
8. Fertilizer	Starter	0	
o. Terminer	Other	0	
9.	Subtotal (sum of line 6, 7, and 8)	207	
10.			
11.	207		

²Use the same units for each line in Table F-4. Include documentation of unit conversion factors used, if any.

Nutrient Management Specifications						
Amount to be applied (lb/ac)	N:0	P ₂ O ₅ :0	K ₂ O:0	Other:0		

Predicted method, form, and timing of application: No application in 2012- Fallow field